Draft version April 28, 2009
Preprint typeset using LaTeX style emulateapj v. 9/08/03

KOZAI CYCLES, TIDAL FRICTION AND THE DYNAMICAL EVOLUTION OF BINARY MINOR PLANETS

Hagai B. Perets[1] and Smadar Naoz[2]

Draft version April 28, 2009

Abstract

In recent years many binary minor planets (BMPs) have been discovered in the Solar system. Many models have been suggested for their formation, but these encounter difficulties explaining their observed characteristics. Here we show that secular perturbations by the Sun (Kozai mechanism) fundamentally change the evolution and the initial distribution of BMPs predicted by such models and lead to unique observational signatures. The Kozai mechanism can lead to a large periodic oscillations in the eccentricity and inclination of highly inclined BMP orbits, where we predict such effects to be observable with current accuracy within a few years (e.g. for the binary asteroid Huenna). In addition, the combined effects of the Kozai mechanism and tidal friction (KCTF) drives BMPs into short period circular orbits. We predict a specific inclination dependent distribution of the separation and eccentricity of BMPs, due to these effects, including a zone of avoidance at the highest inclinations. Specifically the Kozai evolution could explain the recently observed peculiar orbit of the Kuiper belt binary 2001 QW_{322} . Additionally, the KCTF process could lead to BMPs coalescence and serve as an important route for the formation of irregular shaped single minor planets with large axial tilts.

Subject headings:

1. INTRODUCTION

Stable gravitational triple systems require a hierarchical configuration, in which two objects orbit each other in a relatively tight "inner binary", and the third object orbits the binary in a wider "outer binary". Although such triples are stable against disruption, their orbits may change shape and orientation on time scales much longer than their orbital period. In particular, the Kozai-Lidov mechanism(Kozai 1962; Lidov 1962) predicts a secular perturbations of the inner binary orbit. So-called "Kozai oscillations" cause the eccentricity and the inclination to fluctuate. The Kozai mechanism is known to be highly important in the evolution of many triple systems(Kozai 1962; Mazeh & Shaham 1979; Harrington 1968; Kiseleva et al. 1998; Carruba et al. 2002; Nesvorný et al. 2003; Fabrycky & Tremaine 2007). It leads to a large (order unity) periodic oscillations (Kozai cycles) in the eccentricity and inclination of inner binaries with high inclinations with respect to the outer binary orbit (hereafter the relative inclination).

In recent years many binary minor planets [BMPs; both binary asteroids and binary trans-Neptunian objects (TNOs)] have been discovered in the Solar system(Richardson & Walsh 2006). BMPs can be regarded as the inner binary members of a triple system in which the sun is the perturbing companion in the outer orbit. Here we study the importance of such perturbations and show that BMPs are susceptible to Kozai oscillations, which play a major role in their evolution. The combined effects of the Kozai mechanism in addition to tidal friction, (Kozai cycles and tidal friction; KCTF(Mazeh & Shaham 1979; Kiseleva et al. 1998)) which becomes important for BMPs with high eccentricities (induced by the Kozai mechanism), change the orbital parameters of the BMPs. These effects could then erase the observational signatures suggested by the various formation scenarios of BMPs discussed in the literature(Weidenschilling et al. 1989; Weidenschilling 2002; Merline et al. 2002; Goldreich et al. 2002; Funato et al. 2004; Richardson & Walsh 2006; Lee et al. 2007). replacing them with unique and different signatures, that are consistent with current observations.

2. KOZAI OSCILLATIONS

Large Kozai oscillations (see fig. 1; for which we used the Kozai and KCTF evolution code(Fabrycky & Tremaine 2007)) take place when the relative inclination between the inner binary orbit and the outer binary orbit of a triple system is large for initially circular binaries, i.e. $40^\circ \lesssim i \lesssim 140^\circ$ (hereafter Kozai inclinations), with somewhat wider inclination range for initially eccentric BMPs. Systems with Kozai inclinations keep their semi-major axis (SMA) separation, but could be driven into periodic changes of the other orbital parameters of the inner binary. The eccentricity and inclination change in a specific range which depends on these orbital parameters. The maximal eccentricity induced by the Kozai mechanism (assuming small initial eccentricity) is given by

$$e_{max} = \sqrt{1-(5/3)cos^2(i_0)}, \qquad (1)$$

where e_0 and i_0 are the initial eccentricity and inclination, respectively (the more general expressions for the maximal eccentricity and minimal eccentricity (and inclinations) for arbitrary initial paramters have somewhat more lengthy analytic formulation Perets & Naoz 2009). The typical timescale for Kozai oscillations between the limiting values is given by(Kiseleva et al. 1998)

$$P_K = \frac{2P_{out}^2}{3\pi P_{in}} \frac{m_1+m_2+m_3}{m_3} (1-e_{out}^2)^{3/2} \ , \qquad (2)$$

where P_{out} and P_{in} are the orbital periods of the outer and inner binaries in the triple system, respectively;

Electronic address: hagai.perets@weizmann.ac.il
[1] Weizmann Institute of Science, POB 26, Rehovot 76100, Israel
[2] Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel



FIG. 1.— Typical Kozai evolution of three BMPs (note logarithmic time scale on the right). Bold blue curves show the Kozai cycles of the binary TNO similar to 2001 QW$_{322}$ [triangle marks its currently observed(Petit et al. 2008) orbital parameters]. The evolution of 2001 QW$_{322}$ is not affected by tidal friction, and probes a wide range of inclinations and eccentricities. The evolution of the TNO binary 2001 QT$_{297}$ is also shown in dashed line [circle marks its currently observed orbital paramters]. Red curves show the evolution of a hypothetical main belt binary asteroid affected both by the Kozai mechanism and by tidal friction. The BMP periodically evolves into high eccentricities and at later times circularize due to tidal forces and migrates to small SMA separation. Note that the final inclination is typically close to 140° at which eccentricities during the Kozai cycle are highest and tidal friction is therefore more efficient. The small inclination oscillations observed at the final stages are due to tidal torques which become important at this stage(Fabrycky & Tremaine 2007). These could have even larger amplitudes if the spin rate of the BMP members is larger.

$m_1 + m_2$ and m_3 are the masses of the inner binary and the third outer member of the triple (the sun in the case of the BMP-sun triple system considered here), respectively; and e_{out} is the eccentricity of the outer binary.

Many of the BMPs observed in the Solar system are known to have large relative inclinations, making them susceptible to Kozai oscillations (see Perets & Naoz 2009 for an extended discussion of the observed inclinations of BMPs). For such systems the timescale for periodic changes in the absence of any additional forces (such as tidal friction or pericenter precession for non-spherical objects), could be as short as a few thousand years (typical for main belt binary asteroids) and up to 10^7 years for some binary TNOs; much shorter than the lifetime of these systems (see 1 for example). Note that the short, $\sim$ 46 yrs, Kozai timescale for the BMP Huenna [3] may enable a direct observation of the Kozai effect within the next few years, even with current observations accuracy (we find $e_{\max} \simeq 0.23$ and $e_{\min} \simeq 0.15$ for Huenna; i.e. eccentricity change rate of $\sim 3.4 \times 10^{-3}$ yr^{-1}, where current error bars on the eccentricity are of $\pm 6 \times 10^{-3}$; Marchis et al. 2008). The change in eccentricities and inclinations during these timescales could be as large as $\Delta e = e_{\max} - e_{\min} \simeq 0.99 - 0.05 = 0.94$ and $\Delta i = i_{\max} - i_{\min} \simeq 86 - 39 = 47°$ for the known systems (e.g. for the binary TNO 2000 OJ67; based upon the orbital parameters determined by Grundy et al. 2009).



FIG. 2.— The inclination and semi-major axis phase space at which the Kozai, KCTF and Kozai-Hill mechanisms have a dominant role for two observed BMP populations; the binary TNOs (bottom) and the main belt binary asteroids (top). Observed eccentric (×) and circular (filled circles) BMP orbits are shown. BMPs with two inclinations solutions, in which different solutions fall in different regimes are marked by green empty circles (otherwise only one solution is shown, corresponding to better χ^2 score; currently observed BMPs will be discussed extensively in a forthcoming paper; Naoz & Perets, in prep.). BMPs outside the Kozai region (Kozai inclinations; marked by dashed lines between $40° \lesssim i \lesssim 140°$) are negligibly affected by the Kozai evolution. BMPs in the Kozai region can fundamentally change their inclinations and eccentricities and typically have large eccentricities (> 0.05). BMPs in the shaded region are affected by the Kozai mechanism and/or tidal friction and are expected to have relatively small SMA separations and circularized orbits (< 0.05). Stable BMPs should not exist in the the Kozai-Hill region (above the upper arc). The SMA is given in units of the KCTF region baseline, r_c at which BMPs are expected to be circularized.

3. KOZAI CYCLES AND TIDAL FRICTION

The SMA separation of BMPs could evolve due to the Kozai mechanism, when tidal friction effects are accounted for. Such effects become important when the pericenter distance between the BMP members becomes small enough during the Kozai cycles. Orbital energy is then dissipated in each pericenter approach by the tidal friction, leading to the evolution of the BMP to smaller SMA separations and more circular orbits (hereafter Kozai migration or KCTF mechanism(Mazeh & Shaham 1979; Kiseleva et al. 1998) ; see fig. 1). Moreover, such evolution could even lead to mass exchange or coalescence of the BMP members into a single (probably irregularly shaped) minor planet if the separation becomes small enough.

In fig. 2 we show the regions in orbital phase space where the Kozai and KCTF processes become important (only dependence on SMAs and inclinations is shown, eccentricities are assumed to initially be negligible for simplicity). BMPs in the KCTF region, i.e. BMPs with large inclinations and small SMAs such that during their Kozai evolution their pericenter distance becomes small enough (when their eccentricities become large) for tidal friction to dominate their further evolution (see fig. 2) are expected to have small SMA separations and circularized orbits (i.e. low eccentricities). Therefore the KCTF regime is expected to have a zone of avoidance in it's upper region (large SMA separations) from which BMPs are expected to migrate into its lower region (small separations), which is expected to be overpopulated with close BMPs (possibly even contact or coalesced configurations), with low eccentricities. Moreover, the distribution of the inclinations for migrating BMPs in the KCTF could be double peaked, increasing towards 40° and 140°(Fabrycky & Tremaine 2007) , where the pericenter distance is smallest during the Kozai cycle, and therefore tidal friction becomes more dominant and may lock the system from further evolution of the inclination at the final stages of the KCTF evolution (see fig. 1 for such behavior). Note, however, that for fast spinning member of BMPs and/or low mass ratio BMPs, inclinations may continue to fluctuate even at late times of the KCTF evolution, and the inclinations distribution could be strongly affected. In addition, the Kozai mechanism may still excite small eccentricities in BMPs in the KCTF regime. Such excitations could be the source of small but non-zero eccentricities of BMPs in the KCTF regime, such as observed for the binary asteroid Emma and the the Pluto-Charon system in the Kuiper belt (Stern et al. 2003; Marchis et al. 2008). Such excitations are likely to be higher for BMPs with low mass ratios.

---

[3] The inclination of Huenna is outside the Kozai region for initially circular binaries, however eccentric binaries are affected by the Kozai mechanism even at lower eccentricities.

We note that the (usually unknown) asphericity of the binaries components may also affect the BMPs orbital evolution (Chauvineau et al. 1993; Scheeres 1994; Ragozzine & Brown 2009). In addition, minor planets in higher multiplicities (e.g. the triple TNO systems of 2003 EL61) are subjected to mutual forces from additional satellites, not taken into account in our description, which can substantially affect the evolution of the system (Ragozzine & Brown 2009). Such effects are beyond the scope of this paper, where we assume all systems to be binaries with spherical components.

4. KOZAI-HILL INSTABILITY

At the other end, if the apocenter distance of a BMP becomes large enough during the Kozai cycles, the tidal effects of the sun may become dominant. The binary separation then becomes closer to or even larger than the Hill radius of the system; and the BMP is disrupted (hereafter the Kozai-Hill scenario; Perets and Naoz, in preparation). Therefore the Kozai-Hill region (see fig. 2) should be depleted of BMPs. Note that the Hill radius has an inclinations dependence at all inclinations due to the changing strength of the Coriolis force which is proportional to $cos(i)$(Innanen 1980; Hamilton & Burns 1991) . The effect of the Kozai mechanism is an additional effect which effects only the Kozai region. We use an analytic expression for the critical SMA at which a given BMP becomes unstable, which takes into account both these effects (Perets & Naoz 2009). We note that currently observed BMPs indeed have separations that are a few times smaller than the Hill radius, but this could also be related to other processes, such as disruption of the widest binaries through gravitational encounters with other minor or major planets.

5. DISCUSSION

5.1. *KCTF observational signature and BMPs formations scenario*

The Kozai, KCTF and Kozai-Hill processes could appreciably affect the survival of BMPs, their orbits and their orbital phase-space distribution. In fig. 2 we show typical examples of the inclination-SMA separation phase-space in which the different Kozai processes take place. The positions of all BMPs with known inclinations in this phase-space are also shown (detailed discussion of the observed inclinations of BMPs is given elsewhere Perets & Naoz 2009). BMPs in the Kozai region are not susceptible to tidal friction during the Kozai cycle. For such BMPs the eccentricity and inclinations continuously and periodically change between the minimal and maximal Kozai eccentricity (see e.g. fig. 1) and should therefore have a wide distribution of eccentricities which would be very different from the initial conditions set by the formation mechanism of the BMPs. The initial SMA, however, will be conserved in this case, still reflecting the initial conditions.

We note that the current observations (although currently only a small sample exists) are in good agreement with our predictions. All the BMPs in the Kozai region have relatively high eccentricities (none are circular, defined here as $e < 0.05$). The zone of avoidance at the highest inclinations in the KCTF region is empty, and BMPs in the KCTF region tend to be circular and with smaller SMA separations. The large difference between binary TNOs and binary asteroids is likely to be related to different formation mechanisms(Richardson & Walsh 2006) . Note that BMPs formed only at small separations at which tidal friction is important would never be subjected to the pure Kozai mechanism.

The above predictions should be of major concern when trying to constrain BMPs' formation scenarios by observations. Many formations scenarios for BMPs have been described in the literature (see ref. (Richardson & Walsh 2006) for a review) with different predictions for the distribution of BMP SMA separations, eccentricities and inclinations (unfortunately, only few studies explored inclination distribution(Nazzario et al. 2007; Schlichting & Sari 2008) ; given the importance of the Kozai effect we strongly suggest to explore this in future studies). As we have shown the initial conditions as prescribed by different scenarios will be fundamentally changed by the Kozai and KCTF evolution. In order to find evidence for specific formation scenarios of BMPs, it is therefore essential to take the Kozai evolution of the BMPs into account. For example, typical high eccentricity BMPs produced in some scenarios such as the expected $e > 0.8$ in the exchange scenario of Funato et al.(Funato et al. 2004) , or $0.2 \lesssim e \lesssim 0.8$ in the chaos assisted capture scenario(Lee et al. 2007) , could be observed with very low eccentricities, either due to the random phase in which they are observed during their Kozai cycle or due to their KCTF evolution which would have circularized their orbit (e.g. fig. 2). Recently (and after the initial presentation of this manuscript), Petit et al.

(2008) have reported on the peculiar orbit of the Kuiper Belt Binary 2001 QW_{322}, with very large SMA but relatively small eccentricity. Such orbit is difficult to explain through previously suggested BMPs formation scenarios, since a binary with such wide SMA is likely to be produced through the exchange scenario scenario, which would typically produce a highly eccentric binary (Funato et al. 2004). Such orbit, however, is naturally produced through the Kozai mechanism presented here. As shown in fig. 1, the orbit of a binary similar to 2001 QW_{322} could evolve due to the Kozai mechanism from initially high eccentricity and small pericenter distance separation (in fact comparable to the separation of the "regular" Kuiper belt object 2001 QT_{297}) to have a low eccentricity and much larger pericenter distance separation, as observed today. Such a binary could therefore have been formed through a binary-single encounter, and then naturally evolve to its current state through Kozai evolution.

Similarly, typical low eccentricity BMPs, such as might be expected to be produced in the dynamical friction capture scenario(Goldreich et al. 2002) , could be observed with very high eccentricities. Since these effects occur mainly for BMPs in the main Kozai region, we predict an inclination and SMA separation dependence of the eccentricity distribution of BMPs, with a transition between BMPs at the different regimes described in fig. 2. Similarly the SMA separation distribution would also be changed between the different regimes. In fact, only BMPs outside the Kozai region would preserve the eccentricity and SMA separation signature of their formation scenario, whereas in the other regimes these distributions would be dominated by the Kozai mechanism.

In addition to these predictions, we note the importance of environmental effects when combined with Kozai evolution. In some regions of the solar system BMPs were likely to form and/or to evolve for some time in dense environments where they encountered other minor planets (as suggested or required by most BMP formation and evolutionary scenarios). In such an environment, encounters may change the orbital configuration of the binaries in quite a chaotic way(Heggie 1975) especially for the wider binaries that are more susceptible to encounters (larger cross section). The Kozai timescales (Eq. 2) could be shorter than the typical timescale between consequent encounters. In such cases a BMP entering the KCTF regime for the first time following an encounter, could rapidly migrate to smaller SMA separation, thus decreasing its chance for further encounters which could have otherwise potentially export it outside this phase space region. This KCTF region therefore produces a sink in the phase-space distribution and induces a flow from small inclinations to high inclinations in the phase-space distribution of BMPs. We predict that the KCTF mechanism, when at work in a collisional environment, would form a KCTF-collisional signature in which the KCTF and the phase-space regions close to it (i.e. mostly the Kozai region) would be overpopulated relative to the other phase space regions (beside the zone of avoidance which should be depleted of BMPs). The distribution of observed binary TNOs is suggestive of such a signature (fig. 2), with all BMPs at high inclinations, in addition to an empty region at the zone of avoidance, but future observations are required in order to show this at high significance.

5.2. *Binary spin-orbit correlation and irregularly shaped minor planets*

Since KCTF evolution drives BMPs into close configurations, they may become close enough as to evolve into contact configuration or even mergers (similar suggestion was discussed in the context of merger of inner binary in triple stellar systems due to KCTF; Perets & Fabrycky 2009). These contact/merger products may still show a signature of their KCTF formation scenario producing a relative spin-orbit inclination which is likely to be limited to Kozai inclinations. In addition this Kozai-induced merger scenario could naturally explain the existence of large single minor planets with irregular prolonged shapes (e.g. (Hartmann & Cruikshank 1980; Romanishin et al. 2001)) and predict their prolonged axis to be aligned perpendicular to their spin axis (where the latter is expected to have the same biased spin-orbit inclination distribution as the Kozai contact binaries). We note that rotational disruption and direct collisions could also produce irregularly shaped minor planets, but are not expected to produce a bias towards high latitude inclinations. The YORP effect could also form a bias toward specific high inclinations, however this process is efficient only for small asteroids (diameter smaller than 50 km; e.g. Polishook & Brosch 2009, and references there in) , and is not expected to affect TNOs at all. If such mergers products are abundant they could effect the relative spin-orbit inclination distribution, and produce tilt axis distribution which is biased towards high latitudes from the orbital plane. Interestingly, such a bias exists in the pole distribution of large binary asteroids (Kryszczyńska et al. 2007, ; not expected to be affected by YORP).

6. SUMMARY

In this letter we have shown that the secular Kozai perturbations by the sun have a major role in the dynamical evolution of binary minor planets. The Kozai mechanism leads to large periodic oscillations in the eccentricity and inclination of highly inclined BMP orbits and the combined effects of the Kozai mechanism and tidal friction drives BMPs into short period circular orbits. We predict an inclination dependent distribution of the separation and eccentricity of BMPs, due to these effects, including a zone of avoidance at the highest inclinations. Additionally, the KCTF process could lead to BMPs coalescence and serve as an important route for the formation of irregular shaped single minor planets with large axial tilts.

We would like to thank Dan Fabrycky for supplying us with his KCTF evolution code and for his critical review on an early version of this manuscript. We also acknowledge valuable discussions with Darin Ragozzine, Re'em Sari, Tsevi Mazeh and David Polishook. SN acknowledges support from the ISF grant 629/05. We also acknowledge the generous support by the industrial and commercial club of Israel through the Ilan-Ramon scholarship.

REFERENCES

Carruba, V. et al. 2002, Icarus, 158, 434

Chauvineau, B., Farinella, P., & Mignard, F. 1993, Icarus, 105, 370

Fabrycky, D. & Tremaine, S. 2007, ApJ, 669, 1298

Funato, Y. et al. 2004, Nature, 427, 518

Goldreich, P., Lithwick, Y., & Sari, R. 2002, Nature, 420, 643

Grundy, W. M. et al. 2009, Icarus, 200, 627

Hamilton, D. P. & Burns, J. A. 1991, Icarus, 92, 118

Harrington, R. S. 1968, AJ, 73, 190

Hartmann, W. K. & Cruikshank, D. P. 1980, Science, 207, 976

Heggie, D. C. 1975, MNRAS, 173, 729

Innanen, K. A. 1980, AJ, 85, 81

Kiseleva, L. G., Eggleton, P. P., & Mikkola, S. 1998, MNRAS, 300, 292

Kozai, Y. 1962, AJ, 67, 591

Kryszczyńska, A. et al. 2007, Icarus, 192, 223

Lee, E. A., Astakhov, S. A., & Farrelly, D. 2007, MNRAS, 379, 229

Lidov, M. L. 1962, Planetary and Space Science, 9, 719

Marchis, F. et al. 2008, Icarus, 195, 295

Mazeh, T. & Shaham, J. 1979, A& A, 77, 145

Merline, W. J. et al. 2002, Asteroids III, 289

Nazzario, R. C. et al. 2007, Advances in Space Research, 40, 280

Nesvorný, D. et al. 2003, AJ, 126, 398

Perets, H. B. & Fabrycky, D. C. 2009, arXiv:0901.4328

Perets, H. B. & Naoz, S. 2009, In prep.; see arXiv:0809.2095v1

Petit, J.-M. et al. 2008, Science, 322, 432

Polishook, D. & Brosch, N. 2009, Icarus, 199, 319

Ragozzine, D. & Brown, M. E. 2009, arXiv:0903.4213

Richardson, D. C. & Walsh, K. J. 2006, ARAA, 34, 47

Romanishin, W. et al. 2001, PNAS, 98, 11863

Scheeres, D. J. 1994, Icarus, 110, 225

Schlichting, H. E. & Sari, R. 2008, arXiv:0803.0329

Stern, S. A., Bottke, W. F., , & Levison, H. F. 2003, AJ, 125, 902

Weidenschilling, S. J. 2002, Icarus, 160, 212

Weidenschilling, S. J., Paolicchi, P., & Zappala, V. 1989, in Asteroids II, ed. R. P. Binzel, T. Gehrels, & M. S. Matthews (The University of Arizona Press), 643–658